

MINIMAL BOUNDED INDEX SUBGROUP FOR DEPENDENT THEORIES

SAHARON SHELAH

The Hebrew University of Jerusalem
Einstein Institute of Mathematics
Edmond J. Safra Campus, Givat Ram
Jerusalem 91904, Israel

Department of Mathematics
Hill Center-Busch Campus
Rutgers, The State University of New Jersey
110 Frelinghuysen Road
Piscataway, NJ 08854-8019 USA

ABSTRACT. For a dependent theory T , in \mathfrak{C}_T for every type definable group G , the intersection of type definable subgroups with bounded index is a type definable subgroup with bounded index.

The author would like to thank the Israel Science Foundation for partial support of this research (Grant No. 242/03), Publication 876

I would like to thank Alice Leonhardt for the beautiful typing.

First Typed - 05/Mar/28

Latest Revision - 06/Jan/11

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\text{\texttt{TEX}}$

§0 INTRODUCTION

Assume that T is a dependent (complete first order) theory \mathfrak{C} is a $\bar{\kappa}$ -saturated model of T (a monster) G is a type definable (in \mathfrak{C}) group in \mathfrak{C} (of course we consider only types of cardinality $< \bar{\kappa}$).

A type definable subgroup H of G is call bounded if the index $(G : H)$ is $< \bar{\kappa}$. We prove that there is a minimal bounded definable subgroup. The first theorem on this line for T stable is of Baldwin-Saxe [BaSx76].

Recently Hrushovski, Peterzil and Pillay [HPP0x] investigated definable groups, 0-minimality and measure, see also the earlier work on definable subgroups in 0-minimal T in Bezarducci, Otero, Peterzil and Pillay.

Hrushovski in a lecture in the logic seminar in the Hebrew University mentioned that “for every definable group in \mathfrak{C}_T , the intersection of the definable subgroup with bounded index is of bounded index” is proved in [HPP0x] for definable groups with suitable definable measure on them and T dependent; but it is not clear if the existence of definable measure is necessary.

Recent works of the author on dependent theories are [Sh 783] (see §3, §4 on groups) [Sh 863] (e.g. the first order theory of the p-adics is strongly¹ dependent but not strongly² dependent, see end of §1; on strongly² dependent fields see §5) and [Sh:F705]. This work is continued in [Sh:F753] (for G abelian and $\mathbb{L}_{\infty, \bar{\kappa}}$ -definable subgroups).

§1

1.1 Lemma. *For T dependent.*

1) If \circledast below holds then:

- (α) $q(\mathfrak{C})$ is a subgroup of $p(\mathfrak{C})$
- (β) $q(\mathfrak{C})$ is of index $\leq 2^{|T|^{\aleph_0}}$
- (γ) essentially $q(x) \setminus p(x)$ is of cardinality $\leq |T|^{\aleph_0}$ (i.e., for some $q'(x) \subseteq q(x)$ of cardinality $\leq |T|^{\aleph_0}$, $q(x)$ is equivalent to $p(x) \cup q'(x)$).

where

- \circledast (a) $p(x)$ is a type such that $p(\mathfrak{C})$ is a group we call G (under some definable operation xy, x^{-1} and the identity e_G which are constant here; of course all types are of cardinality $< \bar{\kappa}$, \mathfrak{C} is $\bar{\kappa}$ -saturated)
- (b) $q(x) = p(x) \cup \bigcup \{r(x) : r(x) \in \mathbf{R}\}$ where
- (c) $\mathbf{R} = \{r(x) : r(x) \text{ a type such that } (p \cup r)(\mathfrak{C}) \text{ is a sub-group of } p(\mathfrak{C}) \text{ of index } < \bar{\kappa}\}$.

2) There exists some $q' \subseteq q$ over $\text{Dom}(p)$, equivalent to q and such that $|q'| \leq |T| + |\text{Dom}(p)|$. So $(p(\mathfrak{C}) : q(\mathfrak{C})) \leq 2^{|\text{Dom}(p)| + |T|}$.

3) If $r_i(x) \in \mathbf{R}$ for $i < (|T|^{\aleph_0})^+$ then for some $\alpha < (|T|^{\aleph_0})^+$ we have $(p(x) \cup \bigcup \{r_i(x) : i < \alpha\})(\mathfrak{C}) = (p(x) \cup \bigcup \{r_i(x) : i < (|T|^{\aleph_0})^+\})(\mathfrak{C})$.

Proof. 1) Note

- \circledast_1 \mathbf{R} is closed under unions of $< \bar{\kappa}$ hence $q' \subseteq q \wedge |q'| < \bar{\kappa} \Rightarrow q' \in \mathbf{R}$
- \circledast_2 if $r(x) \in \mathbf{R}$, $r'(x) \subseteq r(x)$ is countable then there is a countable $r''(x) \subseteq r(x)$ including $r'(x)$ which belongs to \mathbf{R}
 [Why? Let $r'(x) = \{\varphi_n(x, \bar{a}_n) : n < \omega\}$ (can use $\varphi_n = (x = x)$). Without loss of generality $r(x)$ is closed under conjunctions and also $r'(x)$. Now we choose $\psi_n(x, \bar{b}_n) = \psi_n^1(x, \bar{b}_n^1) \wedge \psi_n^2(x, \bar{b}_n^2)$ with $\psi_n^1(x, \bar{b}_n^1) \in r(x)$, $\psi_n^2(x, \bar{b}_n^2) \in p(x)$ by induction on $n < \omega$ such that $\psi_{n+1}(x, \bar{b}_{n+1}) \wedge \psi_{n+1}(y, \bar{b}_{n+1}) \vdash \psi_n(xy^{-1}, \bar{b}_n) \wedge \varphi_n(x, \bar{a}_n)$. Such formula exists as $(p(x) \cup r(x)) \cup (p(y) \cup r(y)) \vdash \psi_n(xy^{-1}, \bar{b}_n) \wedge \varphi_n(x, \bar{a}_n)$.
 Now $r''(x) = \{\varphi_n(x, \bar{a}_n), \psi_n(x, \bar{b}_n) : n < \omega\}$ is as required.]

Clause (α) is obvious.

Assume toward contradiction that the conclusion (β) fails. So we can choose (c_α, r_α) by induction on $\alpha < (|T|^{\aleph_0})^+$ such that

- ⊗₃ (a) $c_\alpha \in (p(x) \cup \bigcup \{r_\beta : \beta < \alpha\})(\mathfrak{C}) \setminus q(\mathfrak{C})$
- (b) $r_\alpha = \{\psi_n^\alpha(x, \bar{b}_n^\alpha) : n < \omega\} \subseteq q$ and $\bar{b}_n^\alpha \triangleleft \bar{b}_{n+1}^\alpha$
- (c) $\psi_{n+1}^\alpha(x, \bar{b}_{n+1}^\alpha) \vdash \psi_n^\alpha(x, \bar{b}_n^\alpha)$
- (d) $r_\alpha \in \mathbf{R}$
- (e) c_α does not realize r_α in fact $\mathfrak{C} \models \neg \psi_0^\alpha(c_\alpha, \bar{b}_0^\alpha)$.

Without loss of generality

- ⊗₄ $\psi_{n+1}^\alpha(x, \bar{y}_{n+1}^\alpha) \vdash \psi_n^\alpha(x, \bar{y}_n^\alpha)$ and $(\psi_{n+1}^\alpha(x_1, \bar{y}_{n+1}^\alpha) \wedge \psi_{n+1}^\alpha(x_2, \bar{y}_{n+1}^\alpha)) \rightarrow (\psi_{n+1}^\alpha(e, \bar{y}_{n+1}^\alpha) \wedge \psi_{n+1}^\alpha(x_1 x_2^{-1}, \bar{y}_{n+1}^\alpha))$
- ⊗₅ $\psi_n^\alpha(x, \bar{y}_n^\alpha) = \psi_n(x, \bar{y}_n)$ and $\psi_{n+1}(x, \bar{y}_{n+1}) \vdash \psi_n(x, \bar{y}_n)$.
- ⊗₆ $\langle c_\alpha \bar{\mathbf{a}}_\alpha : \alpha < (|T|^{\aleph_0})^+ \text{ is an indiscernible sequence over } \text{Dom}(p) \text{ where } \bar{\mathbf{a}}_\alpha = \bar{b}_0^\alpha \wedge \bar{b}_1^\alpha \wedge \bar{b}_2^\alpha \wedge \dots, \text{ without loss of generality } \bar{b}_n^\alpha = \bar{\mathbf{a}}_\alpha \upharpoonright k_n$
[Why? By Ramsey theorem and compactness.]
- ⊗₇ if $\alpha < \beta < \gamma$ then $c_\alpha c_\beta^{-1} \in r_\gamma(\mathfrak{C})$.
[Why? Without loss of generality γ is infinite, as $(p \cup r_\gamma)(\mathfrak{C})$ is a subgroup of $p(\mathfrak{C})$ of index $< \bar{\kappa}$. If $\gamma \geq \omega$, $\langle c_i : i < \gamma \rangle$ is a sequence of indiscernibles over $\text{Dom}(p) \cup \bar{b}_\gamma$ of elements of $p(\mathfrak{C})$ pairwise non-equivalent modulo $G_\gamma = (p \cup r_\gamma)(\mathfrak{C})$, then extend it to $\langle c_i : i < \bar{\kappa} \rangle$ a sequence of indiscernibles over $\text{Dom}(p) \cup \bar{b}_\gamma$ and arrive at $\alpha < \beta \Rightarrow c_\alpha c_\beta^{-1} \notin G_\gamma \Rightarrow c_\beta c_\alpha^{-1} \notin G_\gamma$ so $\langle c_\alpha G_\gamma : \alpha < \bar{\kappa} \rangle$ pairwise distinct (equivalently $\langle G_\gamma c_\alpha : \alpha < \bar{\kappa} \rangle$ pairwise distinct) contradiction.]
- ⊗₈ $c_\alpha \in r_\beta(\mathfrak{C})$ iff $\alpha \neq \beta$.
[Why? Let

$$c_\alpha^* = c_{2\alpha+1} \cdot (c_{2\alpha})^{-1}$$

$$r_\alpha^* = r_{2\alpha}.$$

So:

- (i) if $\beta < \alpha$, $c_\alpha^* \in (p \cup r_\beta^*)(\mathfrak{C})$ as $c_{2\alpha+1}, c_{2\alpha}$ belong to the subgroup $(p \cup r_{2\beta+1})(\mathfrak{C})$ by clause (a) of ⊗₃
- (ii) if $\beta > \alpha$, c_α^* belongs to $(p \cup r_\beta^*)(\mathfrak{C})$ by ⊗₇
- (iii) if $\beta = \alpha$ then c_α^* does not belong to $(p \cup r_\beta^*)(\mathfrak{C})$ as it is a subgroup, $c_{2\alpha+1}$ belongs to it and $c_{2\alpha}$ does not belong to it by clause (e) of ⊗₃.

Let $\bar{\mathbf{a}}_\alpha^* = \bar{\mathbf{a}}_{2\alpha+1}$, $\bar{b}_n^{\alpha^*} = \bar{b}_n^{2\alpha}$ retaining the same ψ 's. So we have gotten an example as required in \otimes_8 (not losing the other demands).]

\otimes_9 if $d_1, d_2 \in (p \cup r_\alpha)(\mathfrak{C})$ then $d_1 c_\alpha d_2 \notin \varphi_1(\mathfrak{C}, \bar{b}_1^\alpha)$.

[Why? Fix α , if this holds for some $\varphi_n(-, \bar{b}_n^\alpha)$ by indiscernibility renaming the φ_i 's this is O.K. Otherwise for each $n < \omega$ there are $d_1^n, d_2^n \in (p \cup r_\alpha)(\mathfrak{C})$ such that $\mathfrak{C} \models \varphi_n(d_1^n c_\alpha d_2^n, \bar{b}_n^\alpha)$. By compactness for some $d_1^*, d_2^* \in (p \cup r_\alpha)(\mathfrak{C})$ we have $\models \varphi_n[d_1^* c_\alpha d_2^*, \bar{b}_n^\alpha]$ for every $n < \omega$. So $d_1^* c_\alpha d_2^*$ belongs to the subgroup $(p \cup r_\alpha)(\mathfrak{C})$ but also d_1^*, d_2^* belongs to it hence c_α belongs, contradiction.]

\otimes_{10} if $w = \{i_1, \dots, i_n\}$, $i_1 < \dots < i_n < (2^{|T|})^+$, and $d_w := c_{i_1} c_{i_2} \dots c_{i_n}$ then $\models \varphi_1[d_w, \bar{b}_\alpha^1] \Leftrightarrow \alpha \notin w$.

[Why? If $\alpha \in w$ let k be such that $\alpha = i_k$, so $(c_{i_1}, \dots, c_{i_{k-1}}) \in (p \cup r_\alpha)(\mathfrak{C})$ by \otimes_7 and $c_{i_{k+1}} \dots c_{i_n} \in (p \cup r_\alpha)(\mathfrak{C})$ hence $d_w = (c_{i_1} \dots c_{i_k}) c_{i_k} (c_{i_{k+1}} \dots c_{i_n}) \notin (p \cup r_\alpha)(\mathfrak{C})$ by \otimes_9 .

Second, if $\alpha \notin w$ by \otimes_8 as $\{c_{i_\ell} : \ell < n\}$ is included in the subgroup $(p \cup r_\alpha)(\mathfrak{C})$.]

So we get a contradiction to “ T dependent” hence clause (β) holds. Also clause (γ) follows by the following observation:

Observation. If $r(x) \in \mathbf{R}$ and $|r(x)| \leq \theta$ then $(p(\mathfrak{C}) : (p \cup r)(\mathfrak{C})) \leq 2^\theta$ (except finite when θ is finite).

Proof. 1) If θ is finite then by compactness. If σ is infinite then without loss of generality r is closed under conjunctions. Let $r = \{\varphi_i(x, \bar{\mathbf{b}}) : i < \theta\}$, $\bar{\mathbf{b}}$ is possibly infinite.

For each $i < \theta$ let $u \subseteq \text{ord}$ be such that $\bar{\kappa} > |u| > (p(\mathfrak{C}) : (p \cup r)(\mathfrak{C}))$ let $\Gamma_{i,u} = \cup\{p(x_\alpha) : \alpha \in u\} \cup \{\neg\varphi_i(x_\alpha x_\beta^{-1}, \bar{\mathbf{b}}) : \alpha < \beta \text{ from } u\}$. So for some finite $u_i^* \subseteq u$, Γ_{i,u_i^*} is contradictory so Γ_{i,n_i} is contradictory when $n_i = |u_i^*|$. It suffices to use $(2^\theta)^+ \rightarrow (\dots n_i \dots)_{i < \theta}$ (why? let $\langle c_\alpha : \alpha < (2^\theta)^+ \rangle$ exemplify the failure and let $\zeta_{\alpha,\beta} = \text{Min}\{i : \models \neg\varphi_i(c_\alpha c_\beta^{-1}, \bar{\mathbf{b}})\}$).

2) Observe that every automorphism of \mathfrak{C} fixing $\text{Dom}(p)$ maps $p(\mathfrak{C})$ onto itself and therefore maps $q(\mathfrak{C})$ onto itself.

It follows that if $c_1, c_2 \in p(\mathfrak{C})$ are such that $\text{tp}(c_1, \text{Dom}(p)) = \text{tp}(c_2, \text{Dom}(p))$ then $c_1 \in q(\mathfrak{C})$ if and only if $c_2 \in q(\mathfrak{C})$. Let $\mathbf{P} := \{\text{tp}(b/\text{Dom}(p)) | b \in q(\mathfrak{C})\}$, $\mathbf{P}(\mathfrak{C}) := \{r(\mathfrak{C}) : r \subset \mathbf{P}\}$. Then by the above explanation $\mathbf{P}(\mathfrak{C}) \subseteq q(\mathfrak{C})$. By definition $q(\mathfrak{C}) \subseteq p(\mathfrak{C})$ so they are equal. Let $q_{**} = \cap\{r : r \in \mathbf{P}\}$ then $q(\mathfrak{C}) \subseteq q_{**}(\mathfrak{C})$.

If they are equal then we are done. Otherwise take $c_1 \in q_{**}(\mathfrak{C}) \setminus q(\mathfrak{C})$. Without loss of generality let $\psi(x, \bar{d}) \in q$ be such that $\models \neg\psi(c_1, \bar{d})$.

By definition of \mathbf{P} and c_1 , for each $\theta(x, \bar{e}) \in \text{tp}(c_1, \text{Dom}(p))$ there exists some $p_{\theta(x, \bar{e})} \in \mathbf{P}$ such that $\sigma(x, \bar{e}) \in p_{\sigma(x, \bar{e})}$ and therefore some $c_{\sigma(x, \bar{e})} \in q(\mathfrak{C})$ realizes $\theta(x, \bar{e})$. So $\text{tp}(c_1, \text{dom}(p)) \cup q(x)$ is finitely satisfiable and is therefore realized by some c_2 . Thus $\text{tp}(c_1, \text{Dom}(p)) = \text{tp}(c_2, \text{Dom}(p))$, but $c_1 \notin q(\mathfrak{C})$ and $c_2 \in q(\mathfrak{C})$ a contradiction.

3) By the proof of part (1).

$\square_{1.1}$

§2

Claim. *[T dependent] Assume*

- (a) *G is a A^* -definable semi-group with cancellation*
- (b) *$q(x, \bar{a})$ is a type, $q(\mathfrak{C}, \bar{a})$ a sub-semi group.*

Then *we can find q^* and $\langle \bar{a}_i : i < \alpha \rangle$ and B such that*

- (α) $\alpha < (|T|^{\aleph_0})^+$
- (β) $\text{tp}(\bar{a}_i, A^*) = \text{tp}(\bar{a}, A^*)$
- (γ) $q^* = \cup \{q_i(\bar{x}, \bar{a}_i) : i < \alpha\}$
- (δ) $B \subseteq \cap \{q_i(\mathfrak{C}, \bar{a}_i) : i < \alpha\}$ and $|B| \leq |\alpha|$
- (ε) *if \bar{a}' realizes $\text{tp}(\bar{a}, A^*)$ and $B \subseteq q(\mathfrak{C}, \bar{a}')$ then $q^*(\mathfrak{C}) \subseteq q(\mathfrak{C}, \bar{a}')$.*

Proof. We try to choose \bar{a}_α, b_α by induction on $\alpha < (|T|^{\aleph_0})^+$ such that

- (*) (a) \bar{a}_α realizes $\text{tp}(\bar{a}, A^*)$
- (b) $b_\alpha \notin q(\mathfrak{C}, \bar{a}_\alpha)$
- (c) b_α realizes $q(x, \bar{a}_\beta)$ for $\beta < \alpha$
- (d) b_β realizes $q(x, \bar{a}_\alpha)$ for $\beta < \alpha$.

If we succeed we get contradiction as in the proof in §1. If we are stuck at some $\alpha < (|T|^{\aleph_0})^+$ then take $\langle \bar{a}_i : i < \alpha \rangle, B = \{b_i : i < \alpha\}$.

REFERENCES.

- [BaSx76] John T. Baldwin and Jan Saxl. Logical stability in group theory. *J. Austral. Math. Soc. Ser. A*, **21**:267–276, 1976.
- [HPP0x] Ehud Hrushovski, Ya’acov Peterzil, and Anand Pillay. Definable groups, o-minimality and measure. *preprint*.
- [Sh:F753] Saharon Shelah. Definable Groups for Dependent Theories.
- [Sh 783] Saharon Shelah. Dependent first order theories, continued. *Israel Journal of Mathematic*, **accepted**. math.LO/0406440.
- [Sh 863] Saharon Shelah. Strongly dependent theories. *Israel Journal of Mathematics*, **submitted**. math.LO/0504197.
- [Sh:F705] Shelah, Saharon. Representation over orders of elementary classes.